

Annual Male Physical

DOB: Date of last physical:	
Date test last done	Date next due
Colonoscopy	
Prostate exam/ PSA	
Eye exam	
Flu vaccine (Influenza, Trivalent Inactivated, TIV)-typically given once a year in the Fall or Winter. Given intramuscularly. The transnasal vaccine is not recommended for those 50 years of age or older.	
Pneumonia vaccine (Pneumococcal Polysaccharide, PPSV)–recommended for those 65 years or older, if no history or unknown history of receiving it in the past. Second dose given at least five years after first. Recommended for those younger than 65 with certain health conditions.	
Tetanus, Diptheria (td/tdap) vaccine-a routine booster is recommended every ten years after initial series is complete. May be recommended earlier than ten years in certain circumstances. Given intramuscularly.	
Shingles (Zostavax) vaccine–Given once if previously unvaccinated. Given subcutaneously. May receive the vaccination with a history of shingles (VZV) or chicken pox infection.	
Chicken pox vaccine (Varicella, Var)–recommended if no evidence of immunity (previous chicken pox or shingles infection). Given in two subcutaneous doses spaced at least four weeks apart.	
Hepatitis B (Hep B)–recommended for anyone who wishes to be protected from the Hepatitis B virus. Recommended for high risk groups, including those with chronic liver or kidney disease. Given intramuscularly, in a series of three injections spread over at least sixteen weeks.	
Provider/Nurse Signature: Time:	Date: